

Cerulean warbler

Threatened in Quebec – Endangered in Canada



Approx,
10 cm

©Ray Hennessy

CHARACTERISTICS



- Adult male: azure upper body and white underparts, blue-black band across the throat.
- Adult female: turquoise upper body and whitish breast, variegated with yellow.
- Relatively long wings and short tail.



Mid-May



June



Late June to
mid-July




Late August to early
May

HABITAT



- Canopy of large trees (24 to 30 m) in mature, old-growth deciduous forests characterized by sparse understory.
- Requires large woodland patches.
- Nests in the tops of large trees.
- Winters in South America.

DISTRIBUTION IN QUEBEC



Source : CDPNQ

THREATS




- Logging that contributes to fragmentation and reduction of mature forest areas.
- Tree diseases that reduce the number of large trees.
- Conversion of forests for residential, commercial or agricultural development.
- Reduction of insect populations due to the use of pesticides.
- Collision with man-made structures such as communication towers and wind turbines.



Avoid forestry operations during breeding and rearing periods.



Preserve extensive wooded areas with tall trees.



Follow the recommendations in the *Mature deciduous forests* fact sheet.

Illustrations : Sophie-Anne Vachon and Emilie Dumais

For references and to learn more, please visit: foretprivee.ca/biodiversite



Produced by



and



Ce projet a été réalisé avec l'appui financier de :
This project was undertaken with the financial support of:



Environnement et Changement climatique Canada

Environment and Climate Change Canada

